



# Middlesex University Research Repository

An open access repository of

Middlesex University research

<http://eprints.mdx.ac.uk>

Bromley, Tom, Turner, Anthony N., Read, Paul, Lake, Jason, Maloney, Sean, Chavda, Shyam and Bishop, Chris (2018) Effects of a competitive soccer match on jump performance and interlimb asymmetries in elite academy soccer players. Journal of Strength and Conditioning Research . ISSN 1064-8011 (Published online first) (doi:10.1519/JSC.0000000000002951)

Final accepted version (with author's formatting)

This version is available at: <http://eprints.mdx.ac.uk/25399/>

## Copyright:

Middlesex University Research Repository makes the University's research available electronically.

Copyright and moral rights to this work are retained by the author and/or other copyright owners unless otherwise stated. The work is supplied on the understanding that any use for commercial gain is strictly forbidden. A copy may be downloaded for personal, non-commercial, research or study without prior permission and without charge.

Works, including theses and research projects, may not be reproduced in any format or medium, or extensive quotations taken from them, or their content changed in any way, without first obtaining permission in writing from the copyright holder(s). They may not be sold or exploited commercially in any format or medium without the prior written permission of the copyright holder(s).

Full bibliographic details must be given when referring to, or quoting from full items including the author's name, the title of the work, publication details where relevant (place, publisher, date), pagination, and for theses or dissertations the awarding institution, the degree type awarded, and the date of the award.

If you believe that any material held in the repository infringes copyright law, please contact the Repository Team at Middlesex University via the following email address:

[eprints@mdx.ac.uk](mailto:eprints@mdx.ac.uk)

The item will be removed from the repository while any claim is being investigated.

See also repository copyright: re-use policy: <http://eprints.mdx.ac.uk/policies.html#copy>

# THE EFFECTS OF A COMPETITIVE SOCCER MATCH ON JUMP PERFORMANCE AND INTER-LIMB ASYMMETRIES IN ELITE ACADEMY SOCCER PLAYERS

## ABSTRACT

The purpose of the present study was to investigate the effects of a competitive soccer match on jump performance and inter-limb asymmetries over incremental time points during a 72-hour (h) period. Fourteen elite adolescent players from a professional English category three academy performed single leg countermovement jumps (SLCMJ) pre, post, 24, 48, and 72-h post-match on a single force platform. Eccentric impulse, concentric impulse, peak propulsive force, jump height, peak landing force, and landing impulse were monitored throughout. Inter-limb asymmetries were also calculated for each metric as the percentage difference between limbs. Significant negative changes ( $p < 0.05$ ) in jump performance were noted for all metrics at all time points, with the exception of jump height. Inter-limb asymmetries were metric-dependent and showed very large increases, specifically post-match, with a trend to reduce back towards baseline values at the 48-h time point for propulsive-based metrics. Asymmetries for landing metrics did not peak until the 24-h time point and again reduced towards baseline at 48-h. The present study highlights the importance of monitoring distinct jump metrics, as jump height alone was not sensitive enough to show significant changes in jump performance. However, inter-limb asymmetries were sensitive to fatigue with very large increases post-match. More frequent monitoring of asymmetries could enable practitioners to determine whether existing imbalances are also associated with reductions in physical performance or increased injury risk.

**Key Words:** Performance monitoring, imbalances, recovery

## 26 INTRODUCTION

27 During match play, soccer athletes are required to perform repeated high intensity, intermittent, and  
28 multi-directional actions in unpredictable environments. Specifically, jogging, sprinting, jumping, and  
29 changes of direction are common in soccer which frequently require high levels of unilateral force  
30 production (3,32,37). Mohr et al. (33) documented approximately 1300 individual or combinations of  
31 these actions throughout match-play, which have been shown to occur and subsequently change on  
32 average every five seconds. With players consistently required to perform these actions and react to  
33 external stimuli such as opponents and ball trajectory to name just a couple, asymmetrical loading is a  
34 natural consequence; thus, inter-limb asymmetries are likely a by-product of the sport which has been  
35 shown in comparable team sport athletes (23). In addition, the action of kicking is also likely to  
36 contribute to inter-limb differences in soccer players due to the inherent nature of the non-kicking limb  
37 required to stabilize each player (and thus absorb ground reaction force) during the action itself.  
38 Furthermore, inter-limb asymmetries have been negatively associated with sports performance markers  
39 (6,9,25,31), in addition to increased injury risk about the hip (5), knee (14) and ankle (18). Thus,  
40 quantifying and monitoring changes in asymmetry could be deemed important to both maximize  
41 physical performance and reduce potential injury risk.

42 The relationship between fatigue and physical performance has been found increasingly important as  
43 reduced physical performance inclusive of: total distance covered, high speed running, sprint distance,  
44 accelerations, and decelerations (1,45) have also been noted during the latter periods of each half.  
45 Chronic effects of neuromuscular fatigue have also been shown to remain up to 48 hours post-match  
46 (2,27,44). Within the available literature, simulated match protocols have been utilized to induce player  
47 fatigue and monitor both their acute and chronic effects on performance. Significant intra-limb  
48 decreases of 13-15% in functional hamstrings to quadriceps strength ratio (H:Q) have been shown  
49 following the Loughborough intermittent shuttle test and a soccer-specific aerobic field test,  
50 respectively (10,43). Soccer players are generally involved in > 70 matches per season with ~3-6  
51 training sessions every week, leaving little time for recovery (27,41). Thus, the time course of residual

periods where elevated asymmetry may be present also requires examination to inform training prescription and identify vulnerable periods where players may be at a greater risk of injury.

Previous studies have focused on the use of isokinetic dynamometry to measure inter-limb asymmetries (14,16,43). Although useful, isokinetic assessments involve isolated joint actions under the constraints of laboratory conditions and are likely to be time-inefficient in comparison to alternative field-based tests. More recent investigations have highlighted the use of unilateral jump tests to quantify asymmetries in soccer players such as the single leg countermovement jump (SLCMJ) and various hop tests (9). Bishop et al. (9) reported jump height asymmetries of ~12% in youth female players from the SLCMJ, which were double that of any horizontal hop tests. Furthermore, these inter-limb differences were associated with reduced sprint ( $r = 0.49$  to  $0.59$ ) and jump performance ( $r = -0.47$  to  $-0.58$ ) suggesting that players with smaller asymmetries outperformed those with larger differences. However, test protocols in the aforementioned study were performed when players were fresh. At present, the association between side-to-side differences and fatigue in soccer is particularly scarce with studies only using simulated protocols to induce fatigue (10,16,29,43). These protocols exclude the auxiliary actions synonymous with soccer (such as physical duels); thus, not providing a true representation of the demands of competitive match-play.

Therefore, the first aim of the present study was to quantify inter-limb asymmetries in a cohort of elite academy soccer players. The second aim was to determine the effects that a competitive soccer match has on these side-to-side differences. The final aim was to monitor these asymmetries over a 72-h period, which would provide an insight into the relationship between asymmetries, fatigue, and recovery. It was hypothesized that asymmetries would significantly increase post-match with notable reductions seen throughout the 72-h recovery period.

## **METHODS**

### **Experimental Approach to the Problem**

This study examined the acute effects of a single 90-minute soccer match during the in-season period on inter-limb asymmetries in elite male youth soccer players throughout a 72-h time period. Asymmetries were measured using the SLCMJ with all testing conducted on a uniaxial force platform. The SLCMJ had been included in previous strength and conditioning programs bi-weekly for up to six weeks pre-testing, ensuring that all players were fully familiar with the testing protocols. In addition, a complete simulated familiarization session was carried out seven days prior to the experimental trial. Assessments were conducted at scheduled intervals: two hours pre-match, one hour post-match and 24-h, 48-h, and 72-h post-match. To limit external influences, subjects were asked to maintain regular diet and sleeping habits throughout the duration of the study, (details of which were previously provided to players as part of the club's in-house player development program). Within-session reliability was computed three ways at each time point, noting that reliability of some force platform metrics have been affected when athletes are in a fatigued state (21).

### **Subjects**

Fourteen elite adolescent male soccer players (age:  $17.6 \pm 0.5$  yr; body mass:  $63.2 \pm 6.7$  kg; height:  $1.77 \pm 0.8$  m) from a professional English category three academy volunteered to participate in this study. All subjects were regularly completing six hours of technical soccer training and three hours of supplementary strength and conditioning training per week. All subjects had a minimum soccer specific and resistance training age of two years. All participants were free from injury and illness at the time of testing and for the duration of the study period. Parental and participant consent was obtained prior to commencement of the study owing to the participant age and ethical approval was granted from the appropriate institutional review board.

### **Procedures**

*Single Leg Countermovement Jumps (SLCMJ).*

Participants stood on the center of a force plate (400 series performance force plate; Fitness Technology, Australia) operating at 600 Hz, motionless for 2-seconds enabling their system mass to be calculated. Upon instruction, subjects performed a countermovement to a self-selected depth followed immediately by triple extending at the ankles, knees, and hips performing a maximal effort vertical jump. Instructions were to “jump as fast and as high as possible after my countdown”. Subjects were required to keep their hands on hips and legs fully extended at all times during the flight phase of the jump; any deviation from these resulted in a void trial and subsequently retaken after a 30-second rest period. The non-jumping limb was required to remain slightly flexed at the hip and knee so that the foot was hovering approximately parallel to the mid-shin of the jumping limb, with no swinging allowed. This was monitored closely by an accredited strength and conditioning coach to ensure consistency throughout all testing protocols. Prior to the assessment protocol, participants completed a standardized warm up consisting of lower body dynamic stretches (multi-planar lunges, inchworms, ‘world’s greatest stretch’ and squat variations) and practice jumps at 60, 80, and 100% of maximum perceived effort. For data collection, all subjects performed three trials on each limb at each time point, separated by a 30-second rest period between each trial.

Force-time data were analyzed to obtain the dependant variables and manually extracted before being transferred to a personal computer at 600 Hz through USB, which was initially examined through custom made software (Ballistic measurement system, XPV7; Fitness Technology, Australia). The dependant variables for the propulsive phase were: eccentric impulse (the sum of impulse from the end of unweighting period up until the end of the braking phase), concentric impulse (the sum of impulse from the end of the braking phase up until take off), peak force (maximum force obtained during the propulsive phase of the jump), and jump height (jump height was calculated using the velocity at take-off). For the landing phase: landing impulse (the sum of impulse upon landing up until peak landing force) and peak landing force (maximum force obtained during the landing phase of the jump) were all later calculated in Microsoft Excel with force thresholds calculated from body weight  $\pm$  5 standard deviations (SD) (11,17,36).

**Statistical Analysis**

All statistical analysis was completed using SPSS Statistics software (version 21.0; SPSS, Inc., Armonk, NY, USA) with data presented as mean  $\pm$  SD. The normality of data was identified using the Shapiro-Wilk test. Previous research has highlighted reduced consistency and reliability when interpreting athlete data in a fatigued state (12); thus, reliability was computed at each time point. Reliability was calculated for each metric using the coefficient of variation (CV), a two-way random intraclass correlation coefficient (ICC) with absolute agreement and 95% confidence intervals, and the standard error of the measurement (SEM). CV were calculated in Microsoft Excel using the formula  $(SD/average)*100$  and the SEM computed via the formula  $SD*\sqrt{(1-ICC)}$ . ICC's  $\geq 0.70$  and CV  $< 10\%$  were considered acceptable (4,12). A one-way repeated measures ANOVA was used to compare the dependant variables in relation to each time point with statistical significance accepted at  $p < 0.05$  and post-hoc Bonferroni testing was used when differences were identified. Cohen's  $d$  effect sizes (ES) were calculated to determine magnitude of change and interpreted in line with previous suggestions: trivial =  $< 0.35$ ; small =  $0.35-0.80$ ; moderate =  $0.80-1.5$ ; large =  $> 1.5$  (40). Finally, inter-limb asymmetries were quantified using a standard percentage difference method:  $100/(max\ value)*(min\ value)*-1+100$  in line with previous suggestions (8), and the change in asymmetries were reported at each time as a percentage change relative to the baseline value.

## RESULTS

### Within-session Reliability

Upon further analysis illustrated in Table 1, it was found that the majority of metrics demonstrated acceptable reliability and consistency in values. However, jump height and peak landing GRF were found to show notably lower reliability ( $ICC \leq 0.69$ ), in addition to landing impulse and concentric impulse which demonstrated remarkably high variability (CV range = 18 to 30%) across a variety of different time points.

\*\*\* INSERT TABLE 1 ABOUT HERE \*\*\*

### Change in Mean Scores

A representation of all mean data and their subsequent changes in performance for each time point are shown in Table 2. Group means were found to significantly ( $p < 0.05$ ) decrease from baseline at all time points for eccentric impulse, peak propulsive GRF and landing impulse, and significantly increase for peak landing GRF. Significant decreases were also found from post and 24-h to 48-h and 72-h for eccentric impulse and peak propulsive GRF. Further significant decreases were noted between 48-h to 72-h on the left side only for eccentric impulse and propulsive peak GRF. Finally, jump height was only found to be significantly reduced on the left side between post to 24-h and increased between 24-h to 72-h.

\*\*\* INSERT TABLE 2 ABOUT HERE \*\*\*

### Change in Inter-limb Asymmetries

Inter-limb asymmetry values for each time point are shown in Table 3 and Figure 1. SLCMJ mean asymmetries significantly increased ( $p < 0.05$ ) from pre to post and/or 24-h for concentric and eccentric impulse, peak propulsive GRF, and peak landing GRF with small to large ES (0.37-3.15). Significant



reductions in asymmetry were shown from post and 24-h to 48-h and 72-h for eccentric impulse, peak propulsive GRF and peak landing GRF.

\*\*\* INSERT TABLE 3 AND FIGURE 1 ABOUT HERE \*\*\*

**DISCUSSION**

The aim of the present study was to quantify inter-limb asymmetries and jump performance in elite academy soccer players before and after a match at incremental time points. Significant changes in side-to-side differences and jump performance were observed at various time points with a trend for the largest asymmetries and reduction in jump performance evident immediately post-match. As per the hypothesis, inter-limb asymmetries returned to a similar level as the pre-match values at the 48-h time point; however, jump performance for multiple metrics was still significantly reduced. These results indicate that assessing multiple metrics during jump performance may be a more sensitive means of identifying a player's readiness to recover in comparison to asymmetry alone.

Table 1 shows reliability data for CMJ metrics at all time points. The relevance here being that when players have competed (and are in a fatigued state), this may impact the reliability of collected data; thus, understanding its usability was required. Noting that CV values were considered acceptable if < 10%, the only metrics to consistently show acceptable CV were peak force (right), landing impulse (right) and peak landing force (both). Typically, there was a trend for test variability to be lower at baseline prior to the match; thus, the effects of a competitive soccer match may have had a detrimental effect on the reliability of some metrics (e.g., eccentric impulse). However, it is important to highlight that concentric impulse and jump height showed the largest variability (CV range = 18 to 30%); thus, practitioners should be mindful of such metrics if monitoring unilateral jump performance during the recovery period post-matches. Furthermore, when investigating 72-h in particular, it was clear to see that due to the addition of a light tactical training session on the afternoon of 48-h, the reliability of some metrics (such as jump height) were again detrimentally affected. Due to the constraints of congested fixtures which is associated with professional soccer, it was not possible to allow these players three days of recovery. Given the variation in metric reliability across time points, this further supports the notion that in-depth jump analysis is key when interpreting real change (12,20,21).

Table 2 shows the mean scores for all jump metrics. Previously, it has been reported that impairment of neuromuscular function is present up to 72-h post-match, with the 0 to 24-h showing the greatest

reduction in jumping and aerobic performance (12,20,29,42). Results from the present study showed that only concentric impulse and jump height were fully recovered on both limbs at the 48-h time point, with the remaining metrics still exhibiting significant differences from baseline two days post-match. Notably, of all the metrics analysed, eccentric impulse and peak propulsive GRF yielded the greatest significant change in means over the course of 24-h post-match ( $d$  range: post = 5.78-3.40, 24-h = 4.56-2.99). A reduction in eccentric impulse capacity at all time points compared to baseline, likely resulted in athletes altering their jump strategy, allowing time to produce the necessary force without significantly affecting jump height in a fatigued state; thus, warranting the investigation of further metrics other than jump height alone (13,20,21). Furthermore, jump height was shown to have no significant decreases in performance post-match; rather, a small, non-significant increase was actually seen on the left side at this time point. These results perhaps suggest that jump height maybe not be sensitive enough to detect significant immediate change as athletes may mask fatigue by compensating using different strategies (as discussed previously). When considering the landing metrics, landing impulse and peak landing GRF were both sensitive enough to show large significant increases in force-time data at each time point compared to baseline, which has also been shown in comparable research (35). Although challenging to explain with certainty, it seems plausible to suggest that a loss of neuromuscular control may explain why significant increases in landing forces were experienced. The effects of a competitive match clearly resulted in ‘heavier landings’ which represent a serious consideration for practitioners. Given that previous research has identified landing mechanics as being a potential risk factor for injury (15,19,24), practitioners could consider landing-based metrics as useful markers of readiness to train during the recovery process.

Table 3 and Figure 1 show the changing nature of inter-limb asymmetries at each time point. Initially, ECC impulse and peak propulsive GRF showed the highest pre-match leg asymmetries (14.24 and 14.71, respectively). These were shown to significantly elevate immediately post-match increasing 2.25 and 2.16 times greater than baseline with large ES ( $d = 3.15$  and  $2.80$ , respectively), then plateauing over the succeeding 24-h before returning to near baseline 48-h. Notably, jump height was found to display the greatest change in asymmetry, increasing immediately post-match by 3.7 times greater than

pre-match, representing a moderate ES ( $d = 1.18$ ). This is particularly striking as this metric showed just a 4.65% asymmetry value pre-match, which can be considered small (7). The relevance being that this has the potential to mislead practitioners unless monitored throughout the time course of 24 to 48-h. Furthermore, given that jump height also showed high variability (CV range = 18 to 30%), this further highlights the need to monitor multiple metrics during jump profiling, and not rely solely on outcome measures. Interestingly, landing metrics were found to take longer to exhibit their peak leg asymmetries; exponentially worsening until 24-h where both metrics climbed to 2.24 and 2.11 times greater than baseline displaying large to moderate ES for landing impulse ( $d = 1.7$ ) and peak landing GRF ( $d = 1.38$ ), respectfully.

When interpreting asymmetries in a more general manner, previous research has highlighted the test and metric-specific nature of asymmetries (9,14,23,26,28,39), and to the authors' knowledge, this is the first study to examine the effects of competition on inter-limb asymmetries in an elite soccer population. Previous research has suggested that 15% might be a threshold where the risk of injury increases (26); however, more recent suggestions advocate 10% as a target to aim for (28,34). When this conflicting evidence is considered and the test-specific nature of asymmetries is deliberated, it seems prudent to suggest that multiple metrics be considered to further our understanding of how inter-limb asymmetries interact with measures of physical performance and injury risk. In the present study, the largest inter-limb differences were noted for peak propulsive force and eccentric impulse (Figure 1). Noting that these metrics can only be obtained from a force platform and showed a trend to exhibit substantially greater differences than the outcome measure of jump height, it is suggested (where possible) that the monitoring of inter-limb differences is conducted using force platforms. As a final point, it is worth noting that it is unlikely that elite soccer players will be granted three days of recovery after matches. With a light tactical training session prescribed after the 48-h time point, this may explain the subsequent increase in asymmetries at 72-h. The results highlight the importance of frequent asymmetry monitoring, which has been emphasized in previous literature (7,35).

## **PRACTICAL APPLICATIONS**

The findings of the present study highlight the changing nature of jump performance and inter-limb asymmetries after a competitive soccer match. Strength and conditioning coaches should consider using the unilateral CMJ in addition to or in place of the more commonly accepted neuromuscular fatigue monitoring method of bilateral CMJ. Noting that many of the actions in soccer occur unilaterally (such as sprinting, changing direction, and kicking), it seems logical to suggest that unilateral jump profiling serves as an ecologically valid means for soccer athletes. In addition, given the previously reported associations between inter-limb asymmetries and reduced physical performance and injury risk, frequent monitoring of side-to-side differences may provide practitioners with a true picture of the interaction between asymmetry and performance or injury risk. The relevance here being that literature pertaining to the longitudinal tracking of asymmetry is scarce. Finally, with leg asymmetries highlighted up to 72-h, this insight can further objectively enlighten the coaching staff on player welfare and subsequently inform their approach to adapting training loads on an individual level; thus, improving training quality, player readiness and ultimately, match performance.

## REFERENCES

1. Akenhead R, Hayes PR, Thompson KG, and French D. Diminutions of acceleration and deceleration output during professional football match play. *J Sci Med Sport* 16: 556-561, 2013.
2. Ascensão A, Rebelo A, Oliveira E, Marques F, Pereira L, and Magalhães J. Biochemical impact of a soccer match - analysis of oxidative stress and muscle damage markers throughout recovery. *Clin Biochem* 41: 841-851, 2008.
3. Bangsbo J, Mohr M, and Krstrup P. Physical and metabolic demands of training and match-play in the elite football player. *J Sports Sci* 24: 665-674, 2006.
4. Baumgartner TA, and Chung H. Confidence Limits for Intraclass Reliability Coefficients. *Meas Phys Edu Exerc Sci* 5: 179-188, 2001.
5. Belhaj K, Meftah S, Mahir L, Lmidmani F, and Elfatimi A. Isokinetic imbalance of adductor–abductor hip muscles in professional soccer players with chronic adductor-related groin pain. *Eur J Sport Sci* 16: 1226-1231, 2016.
6. Bell DR, Sanfilippo JL, Binkley N, and Heiderscheit BC. Lean mass asymmetry influences force and power asymmetry during jumping in collegiate athletes. *J Strength Cond Res* 28: 884-891, 2014.
7. Bishop C, Turner A, and Read P. Effects of inter-limb asymmetries on physical and sports performance: A systematic review. *J Sports Sci* 36: 1135-1144, 2018.
8. Bishop C, Read P, Lake J, Chavda S, Turner A. Inter-limb asymmetries: Understanding how to calculate differences from bilateral and unilateral tests. *Strength Cond J* (Published ahead of print).
9. Bishop C, Read P, McCubbine J, and Turner A. Vertical and Horizontal Asymmetries are Related to Slower Sprinting and Jump Performance in Elite Youth Female Soccer Players: *J Strength Cond Res* (Published ahead of print).
10. Camarda SR, and Denadai BS. Does muscle imbalance affect fatigue after soccer specific intermittent protocol? *J Sci Med Sport* 15: 355-360, 2012.
11. Chavda S, Bromley T, Jarvis P, Williams S, Bishop C, Turner A, Lake J, and Mundy P. Force-time characteristics of the countermovement jump: Analyzing the curve in excel. *Strength Cond J* 40: 67-77, 2018.
12. Cormack SJ, Newton RU, McGuigan MR, and Cormie P. Neuromuscular and endocrine responses of elite players during an Australian rules football season. *Int J Sports Physiol Perform* 3: 439-453, 2008.
13. Cormie P, McBride JM, and McCaulley GO. Power-Time, Force-Time, and Velocity-Time Curve analysis of the countermovement jump: Impact of training: *J Strength Cond Res* 23: 177-186, 2009.

14. Croisier J-L, Ganteaume S, Binet J, Genty M, and Ferret J-M. Strength imbalances and prevention of hamstring injury in professional soccer players: A prospective study. *Am J Sports Med* 36: 1469-1475, 2008.
15. Dai B, Mao D, Garrett WE, and Yu B. Anterior cruciate ligament injuries in soccer: Loading mechanisms, risk factors, and prevention programs. *J Sport Health Sci* 3: 299-306, 2014.
16. Delextrat A, Baker J, Cohen DD, and Clarke ND. Effect of a simulated soccer match on the functional hamstrings-to-quadriceps ratio in amateur female players: H:Q ratio in female soccer players. *Scand J Med Sci Sports* 23: 478-486, 2013.
17. Dos'Santos T, Jones PA, Comfort P, and Thomas C. Effect of different onset thresholds on isometric mid-thigh pull force-time variables. *J Strength Cond Res* 31: 3463-3473, 2017.
18. Fousekis K, Tsepis E, and Vagenas G. Intrinsic risk factors of non-contact ankle sprains in soccer: A prospective study on 100 professional players. *Am J Sports Med* 40: 1842-1850, 2012.
19. Gardiner JC, Weiss JA, and Rosenberg TD. Strain in the human medial collateral ligament during valgus loading of the knee. *Clin Orthop Rel Res* 391: 266-274, 2001.
20. Gathercole R, Sporer B, Stellingwerff T, and Sleivert G. Alternative countermovement jump analysis to quantify acute neuromuscular fatigue. *Int J Sports Physiol Perform* 10: 84-92, 2015.
21. Gathercole RJ, Stellingwerff T, and Sporer BC. Effect of acute fatigue and training adaptation on countermovement jump performance in elite snowboard cross athletes: *J Strength Cond Res* 29: 37-46, 2015.
22. Gibson NE, Boyd AJ, and Murray AM. Countermovement jump is not affected during final competition preparation periods in elite rugby sevens players. *J Strength Cond Res* 30: 777-783, 2016.
23. Hart NH, Nimphius S, Weber J, Spiteri T, Rantalainen T, Dobbin M, and Newton RU. Musculoskeletal asymmetry in football athletes: A product of limb function over time. *Med Sci Sports Exerc* 48: 1379-1387, 2016.
24. Hewett TE, Myer GD, and Ford KR. Decrease in neuromuscular control about the knee with maturation in female athletes. *J Bone Joint Surg* 86: 1601-1608, 2004.
25. Hoffman JR, Ratamess NA, Klatt M, Faigenbaum AD, and Kang J. Do bilateral power deficits influence direction-specific movement patterns? *Res Sports Med* 15: 125-132, 2007.
26. Impellizzeri FM, Rampinini E, Maffiuletti N, and Marcora SM. A vertical jump force test for assessing bilateral strength asymmetry in athletes. *Med Sci Sports Exerc* 39: 2044-2050, 2007.
27. Ispirlidis I, Fatouros IG, Jamurtas AZ, Nikolaidis MG, Michailidis I, Douroudos I, Konstantinos M, Athanasios C, Elias K, Ioannis K, Vassilios K, and Kiriakos T. Time-course of changes in inflammatory and performance responses following a soccer game. *Clin J Sport Med* 18: 423-431, 2008.

28. Kyritsis P, Bahr R, Landreau P, Miladi R, and Witvrouw E. Likelihood of ACL graft rupture: Not meeting six clinical discharge criteria before return to sport is associated with a four times greater risk of rupture. *Brit J Sports Med* 50: 946–951, 2016.
29. Magalhães J, Rebelo A, Oliveira E, Silva JR, Marques F, and Ascensão A. Impact of Loughborough Intermittent Shuttle Test versus soccer match on physiological, biochemical and neuromuscular parameters. *Eur J Appl Physiol* 108: 39-48, 2010.
30. Malone JJ, Murtagh CF, Morgans R, Burgess DJ, Morton JP, and Drust B. Countermovement jump performance is not affected during an in-season training microcycle in elite youth soccer players: *J Strength Cond Res* 29: 752-757, 2015.
31. Menzel H-J, Chagas MH, Szmuchrowski LA, Araujo S, de Andrade A, and de Jesus-Moraleida FR. Analysis of lower limb asymmetries by isokinetic and vertical jump tests in soccer players. *J Strength Cond Res* 27: 1370-1377, 2013.
32. Mohr M, Krstrup P, and Bangsbo J. Fatigue in soccer: A brief review. *J Sports Sci* 23: 593-599, 2005.
33. Mohr M, Krstrup P, and Bangsbo J. Match performance of high-standard soccer players with special reference to development of fatigue. *J Sports Sci* 21: 519-528, 2003.
34. Myer GD, Paterno MV, Ford KR, Quatman CE, and Hewett TE. Rehabilitation after anterior cruciate ligament reconstruction: Criteria-based progression through the return-to-sport phase. *J Orthop Sports Phys Ther* 36: 385-402, 2006.
35. Oliver J, Armstrong N, and Williams C. Changes in jump performance and muscle activity following soccer-specific exercise. *J Sports Sci* 26: 141-148, 2008.
36. Owen NJ, Watkins J, Kilduff LP, Bevan HR, and Bennett MA. Development of a criterion method to determine peak mechanical power output in a countermovement jump. *J Strength Cond Res* 28: 1552-1558, 2014.
37. Rampinini E, Coutts A, Castagna C, Sassi R, and Impellizzeri F. Variation in top level soccer match performance. *Int J Sports Med* 28: 1018-1024, 2007.
38. Read P, Oliver J, De Ste Croix M, Myer G, and Lloyd R. Consistency of field-based measures of neuromuscular control using force-plate diagnostics in elite male youth soccer players. *J Strength Cond Res* 30: 3304-3311, 2016.
39. Read P, Oliver J, De Ste Croix M, Myer G, and Lloyd R. The effects of maturation on measures of asymmetry during neuromuscular control tests in elite male youth soccer players. *Ped Exerc Sci* 30: 168-175, 2017.
40. Rhea MR. Determining the magnitude of treatment effects in strength training research through the use of the effect size. *J Strength Cond Res* 18: 918-920, 2004.



- 404 41. Scott BR, Lockie RG, Knight TJ, Clark AC, and Janse de Jonge X. A comparison of methods  
405 to quantify the in-season training load of professional soccer players. *Int J Sports Physiol*  
406 *Perform* 8: 195-202, 2013.
- 407 42. Silva JR, Rebelo A, Marques F, Pereira L, Seabra A, Ascensão A, and Magalhaes J.  
408 Biochemical impact of soccer: An analysis of hormonal, muscle damage, and redox markers  
409 during the season. *Appl Physiol Nut Metabol* 39: 432-438, 2014.
- 410 43. Small K, McNaughton L, Greig M, and Lovell R. The effects of multidirectional soccer-specific  
411 fatigue on markers of hamstring injury risk. *J Sci Med Sport* 13: 120-125, 2010.
- 412 44. Thomas K, Dent J, Howatson G, and Goodall S. Etiology and recovery of neuromuscular  
413 fatigue following simulated soccer match-play. *Med Sci Sports Exerc* 45: 955-964, 2017.
- 414 45. Weston M, Batterham AM, Castagna C, Portas MD, Barnes C, Harley J, and Lovell R.  
415 Reduction in physical match performance at the start of the second half in elite soccer. *Int J*  
416 *Sports Physiol Perform* 6: 174-182, 2011.

417 Table 1: Within-session reliability data for unilateral countermovement jump metrics at pre, post, 24, 48 and 72 hours post-match.

<b>CMJ Metric</b>	<b>Pre</b>	<b>Post</b>	<b>24-h</b>	<b>48-h</b>	<b>72-h</b>
Eccentric impulse (R)					
CV (%)	6.61	9.21	11.78	12.12	8.50
ICC (95% CI)	0.95 (0.82-0.99)	0.96 (0.87-0.99)	0.98 (0.97-0.99)	0.95 (0.34-0.99)	0.90 (0.45-0.97)
SEM	2.74	2.43	2.25	4.06	3.87
Eccentric impulse (L)					
CV (%)	8.76	10.59	13.12	11.94	11.39
ICC (95% CI)	0.93 (0.77-0.98)	0.89 (0.65-0.96)	0.97 (0.89-0.99)	0.84 (0.69-0.96)	0.97 (0.80-0.99)
SEM	3.51	3.22	2.24	6.18	2.32
Concentric impulse (R)					
CV (%)	18.25	17.99	20.95	21.04	21.18
ICC (95% CI)	0.77 (0.42-0.92)	0.76 (0.42-0.92)	0.89 (0.69-0.97)	0.74 (0.38-0.91)	0.71 (0.32-0.89)
SEM	13.92	10.94	9.34	15.20	15.21
Concentric impulse (L)					
CV (%)	20.29	20.73	22.48	23.97	20.73
ICC (95% CI)	0.85 (0.55-0.95)	0.77 (0.29-0.92)	0.75 (0.27-0.92)	0.84 (0.52-0.95)	0.77 (0.27-0.92)
SEM	11.43	11.29	12.47	12.41	11.22
Peak force (R)					
CV (%)	5.52	8.15	9.67	8.35	7.91
ICC (95% CI)	0.80 (0.58-0.95)	0.92 (0.77-0.97)	0.95 (0.87-0.99)	0.56 (0.11-0.83)	0.85 (0.60-0.95)
SEM	41.05	29.48	26.57	78.83	40.04
Peak force (L)					
CV (%)	10.38	10.68	14.33	12.28	11.55
ICC (95% CI)	0.89 (0.86-0.99)	0.95 (0.84-0.98)	0.92 (0.78-0.99)	0.90 (0.78-0.99)	0.99 (0.96-0.99)
SEM	33.69	21.43	35.70	46.13	12.75
Jump height (R)					
CV (%)	19.51	29.29	19.68	22.19	23.48
ICC (95% CI)	0.89 (0.71-0.96)	0.69 (0.27-0.89)	0.92 (0.78-0.97)	0.89 (0.71-0.96)	0.77 (0.43-0.92)
SEM	0.01	0.03	0.01	0.01	0.02

## Monitoring Asymmetries and Jump Performance

Jump height (L)					
CV (%)	22.15	30.00	25.80	24.23	30.00
ICC (95% CI)	0.85 (0.25-0.96)	0.60 (0.20-0.86)	0.86 (0.60-0.95)	0.72 (0.31-0.90)	0.57 (0.17-0.85)
SEM	0.02	0.04	0.02	0.02	0.04
Landing impulse (R)					
CV (%)	8.96	8.13	7.64	7.05	8.71
ICC (95% CI)	0.68 (0.28-0.88)	0.63 (0.20-0.86)	0.57 (0.11-0.84)	0.58 (0.12-0.84)	0.65 (0.23-0.87)
SEM	1.41	1.69	1.72	1.42	1.72
Landing impulse (L)					
CV (%)	8.96	8.13	16.18	14.81	15.00
ICC (95% CI)	0.81 (0.52-0.93)	0.84 (0.59-0.95)	0.70 (0.32-0.89)	0.81 (0.53-0.93)	0.81 (0.54-0.93)
SEM	1.09	1.10	2.79	1.76	1.93
Peak landing force (R)					
CV (%)	7.48	7.63	7.93	8.86	7.80
ICC (95% CI)	0.90 (0.73-0.97)	0.93 (0.78-0.98)	0.89 (0.70-0.96)	0.88 (0.67-0.96)	0.91 (0.75-0.97)
SEM	62.36	64.96	86.78	91.96	74.32
Peak landing force (L)					
CV (%)	7.48	7.63	8.63	5.72	5.86
ICC (95% CI)	0.74 (0.25-0.94)	0.71 (0.32-0.89)	0.83 (0.26-0.98)	0.64 (0.21-0.87)	0.69 (0.27-0.90)
SEM	101.77	134.15	134.76	109.92	119.97

CMJ = countermovement jump; R = Right; L = Left; CV = coefficient of variation; ICC = intraclass correlation coefficient; CI = confidence interval; SEM = standard error of the measurement.

418

419

420

421

422

423 Table 2: Mean unilateral countermovement jump data  $\pm$  standard deviations for pre, post, 24, 48 and 72 hours post-match.

CMJ Metric	Pre	Post	24-h	48-h	72-h
ECC impulse-R (Ns)	173.68 $\pm$ 11.67 <sup>bcde</sup>	135.54 $\pm$ 12.48 <sup>ac</sup>	134.96 $\pm$ 15.90 <sup>abd</sup>	152.88 $\pm$ 18.52 <sup>abc</sup>	141.24 $\pm$ 12.00 <sup>a</sup>
ECC impulse-L (Ns)	135.54 $\pm$ 13.28 <sup>bce</sup>	92.17 $\pm$ 9.76 <sup>ade</sup>	95.56 $\pm$ 12.53 <sup>ade</sup>	130.97 $\pm$ 15.64 <sup>bce</sup>	110.52 $\pm$ 12.59 <sup>abcd</sup>
CON impulse-R (Ns)	157.64 $\pm$ 28.77 <sup>b</sup>	124.96 $\pm$ 22.47 <sup>a</sup>	135.11 $\pm$ 28.30	141.17 $\pm$ 29.70	132.19 $\pm$ 28.00 <sup>a</sup>
CON impulse-L (Ns)	147.49 $\pm$ 29.92 <sup>bce</sup>	112.35 $\pm$ 23.29 <sup>a</sup>	110.51 $\pm$ 24.84 <sup>a</sup>	129.87 $\pm$ 31.13	112.35 $\pm$ 23.29 <sup>a</sup>
Peak force-R (N)	1679.1 $\pm$ 92.73 <sup>bcde</sup>	1278.52 $\pm$ 104.23 <sup>ade</sup>	1280.76 $\pm$ 123.89 <sup>ade</sup>	1428.09 $\pm$ 119.25 <sup>abc</sup>	1325.51 $\pm$ 104.79 <sup>abc</sup>
Peak force-L (N)	1432.16 $\pm$ 100.66 <sup>bcde</sup>	871.36 $\pm$ 93.08 <sup>ade</sup>	903.53 $\pm$ 129.51 <sup>ade</sup>	1206.53 $\pm$ 148.10 <sup>abce</sup>	1052.69 $\pm$ 121.58 <sup>abcd</sup>
Jump height-R (m)	0.19 $\pm$ 0.04	0.19 $\pm$ 0.06	0.16 $\pm$ 0.03	0.16 $\pm$ 0.04	0.17 $\pm$ 0.04
Jump height-L (m)	0.19 $\pm$ 0.07	0.20 $\pm$ 0.06 <sup>c</sup>	0.16 $\pm$ 0.04 <sup>be</sup>	0.17 $\pm$ 0.04	0.20 $\pm$ 0.06 <sup>c</sup>
Landing impulse-R (Ns)	29.38 $\pm$ 2.22 <sup>bc</sup>	31.18 $\pm$ 2.15 <sup>ae</sup>	31.68 $\pm$ 5.13 <sup>ae</sup>	30.27 $\pm$ 4.04 <sup>c</sup>	29.57 $\pm$ 4.44 <sup>bc</sup>
Landing impulse-L (Ns)	27.83 $\pm$ 2.49 <sup>bce</sup>	34.08 $\pm$ 2.77 <sup>a</sup>	36.20 $\pm$ 2.61 <sup>abd</sup>	31.14 <sup>abc</sup>	31.33 $\pm$ 2.90 <sup>a</sup>
Peak landing force-R (N)	2662.81 $\pm$ 199.21 <sup>bcde</sup>	2963.55 $\pm$ 212.91 <sup>ace</sup>	3271.16 $\pm$ 259.30 <sup>abd</sup>	2983.55 $\pm$ 264.37 <sup>ace</sup>	3195.73 $\pm$ 249.12 <sup>abd</sup>
Peak landing force-L (N)	2811.71 $\pm$ 179.22 <sup>bcde</sup>	3263.55 $\pm$ 249.10 <sup>ace</sup>	3823.25 $\pm$ 329.77 <sup>abd</sup>	3217.99 $\pm$ 184.23 <sup>ace</sup>	3703.72 $\pm$ 216.87 <sup>abd</sup>
CMJ = countermovement jump; ECC = eccentric; CON = concentric; R = Right; L = Left; <sup>a</sup> = significantly different from pre-match value; <sup>b</sup> = significantly different from post-match value; <sup>c</sup> = significantly different from 24-h match value; <sup>d</sup> = significantly different from 48-h match value; <sup>e</sup> = significantly different from 72-h match value.					

Table 3: Inter-limb asymmetry values (reported as percentages) for unilateral countermovement jump data pre, post, 24, 48 and 72 hours post-match and Cohen's *d* effect sizes reported relative to pre-match values.

Asymmetry %	Pre	Post	24-h	48-h	72-h
ECC impulse	14.24 <sup>bce</sup>	32.00 <sup>ade</sup>	29.20 <sup>ade</sup>	14.33 <sup>bc</sup>	21.75 <sup>abc</sup>
Effect size		3.15	2.80	0.01	1.36
CON impulse	7.73 <sup>c</sup>	10.50	18.50 <sup>a</sup>	9.88	15.19
Effect size		0.31	1.02	0.30	0.84
Peak force	14.71 <sup>bc</sup>	31.85 <sup>ade</sup>	29.45 <sup>ade</sup>	15.51 <sup>bc</sup>	20.58 <sup>bc</sup>
Effect size		2.80	2.23	0.12	0.96
Jump height	4.65 <sup>e</sup>	17.22	6.52 <sup>e</sup>	5.47	20.49 <sup>ac</sup>
Effect size		1.18	0.50	0.23	2.05
Peak landing impulse	5.89 <sup>c</sup>	8.51 <sup>cde</sup>	13.14 <sup>abde</sup>	4.03 <sup>bc</sup>	5.61 <sup>bc</sup>
Effect size		0.62	1.71	-0.47	-0.07
Peak landing force	7.22 <sup>ce</sup>	9.13 <sup>ce</sup>	15.22 <sup>abd</sup>	7.29 <sup>ce</sup>	13.72 <sup>abd</sup>
Effect size		0.32	1.38	0.01	1.19
ECC = eccentric; CON = concentric; <sup>a</sup> = significantly different from pre-match value; <sup>b</sup> = significantly different from post-match value; <sup>c</sup> = significantly different from 24-h match value; <sup>d</sup> = significantly different from 48-h match value; <sup>e</sup> = significantly different from 72-h match value.					

## Monitoring Asymmetries and Jump Performance




Figure 1: Inter-limb asymmetry values and standard deviations (error bars) for SLCMJ metrics at pre, post, 24, 48 and 72 hours post-match.